

Effects of Leaf Epidermis Peeling on the CO₂ Exchange Rate and Chlorophyll Fluorescence Quenching, and Estimation of Photorespiration Rate from Electron Transport in Mungbean (*Vigna radiata* (L.) Wilczek) Leaves.

Tokuda, Shinji

Laboratory of Plant Production Physiology, Division of Bioresource and Bioenvironmental Sciences, Graduate School, Kyushu University

Kubota, Fumitake

Laboratory of Plant Production Physiology, Division of Bioresource and Bioenvironmental Sciences, Graduate School, Kyushu University

Hirao, Kenji

Laboratory of Plant Production Physiology, Division of Bioresource and Bioenvironmental Sciences, Graduate School, Kyushu University

Saitou, Kazuyuki

Laboratory of Plant Production Physiology, Division of Bioresource and Bioenvironmental Sciences, Graduate School, Kyushu University

<https://doi.org/10.5109/24273>

出版情報：九州大学大学院農学研究院紀要. 43 (3/4), pp.293-302, 1999-02. 九州大学農学部
バージョン：
権利関係：



Effects of Leaf Epidermis Peeling on the CO₂ Exchange Rate and Chlorophyll Fluorescence Quenching, and Estimation of Photorespiration Rate from Electron Transport in Mungbean (*Vigna radiata* (L.) Wilczek) Leaves.

Shinji Tokuda, Fumitake Kubota, Kenji Hirao and Kazuyuki Saitou

Laboratory of Plant Production Physiology, Division of Bioresource and
Bioenvironmental Sciences, Graduate School, Kyushu University,
Fukuoka 812–8581, Japan

(Received October 1, 1998 and accepted November 6, 1998)

CO₂ gas exchange rates (CER) in a peeled leaf (PL) and nonpeeled leaf (NL) of mungbean (*Vigna radiata* (L.) Wilczek) were measured in a saturating light intensity to identify their responses to CO₂ concentrations ([CO₂]) and leaf temperatures. The CER of PL was regarded as an indicator for the photosynthetic potential in mesophyll. When measured at 35°C, CER was enhanced by peeling the epidermis, whereas it was less effective at 25°C and almost not effective at 15°C. In [CO₂]-CER response curves, CER of PL had a highest plateau in 500 to 1,000 μmol mol⁻¹ [CO₂] at 35°C, whereas at both 25 and 15°C CER showed a peak in 500 μmol mol⁻¹ [CO₂]. When measured in the air of 20% O₂ concentration ([O₂]) in a light intensity of 300 μmol m⁻² s⁻¹ photosynthetically photon flux density (PPFD) at 25°C, the CO₂ quantum yield (Φ_{CO_2}) in PL was higher than that in NL, whereas the photosystem II (PSII) quantum yield (Φ_2) was a little lower in PL. The number (k) of electron equivalents required to fix 1 mol CO₂ was greatly increased in NL due to the increased photorespiration. In 2% [O₂] k of PL was 4.62, which was a little larger than a theoretical minimum (k=4). From the relationships between k, CER and adenosine 5'-triphosphate (ATP) required for photosynthesis, the rates of photorespiration were estimated in different environments.

INTRODUCTION

Mungbean, *Vigna radiata* (L.) Wilczek, is positioned as one of the important protein providing foodstuffs in the tropical and subtropical countries. Many of the studies dealing with mungbean have been conducted mainly on the productivity of field-grown cultivars, but little information is available for physiological aspects in photosynthesis. Accumulating data of the photosynthetic characteristics is necessary as basic information for genetic and physiological improvement of production.

The stomatal resistance is one of the restricting factors for gas exchange, and by removing the resistance the mesophyll activity in photosynthesis can be determined. Experimentally this activity is known by measuring CO₂ exchange rate (CER) of a leaf the epidermis of which has been peeled (Kubota *et al.*, 1991 and 1992). The investigation of mesophyll activity controlled by environmental factors is expected to offer many suggestions for understanding the photosynthetic characteristics of this crop. To clarify the responses of the mesophyll activity to CO₂ concentration ([CO₂]) and leaf temperature is one of the subjects in this study.

Photosynthesis consists of two categories in energy balance, energy production by

photosystem and its consumption by CO₂ assimilation, and the situation of electron transport from photosystem to CO₂ assimilation site is diagnosed by the chlorophyll fluorescence quenching analysis (Schreiber *et al.*, 1986, 1998). When peeled leaves are used for the diagnosis, the relationship between CO₂ assimilation and electron transport should be more clearly interpreted.

The relationships between photosynthesis and chlorophyll fluorescence quenching in leaves were discussed by Genty *et al.* (1989), Marco *et al.* (1990), Krall *et al.* (1990, 1991a, 1991b) and Harbinson *et al.* (1990). In our study the effect of peeling the epidermis or removing the stomatal resistance on CER, photorespiration and electron transport in photosystem II (PSII) were investigated in relation to leaf temperature, [CO₂] and O₂ concentration ([O₂]). On the basis of CER and electron transport, photorespiration rates were estimated in different environments, and the role of photorespiration and its usefulness as a selection criterion in mungbean breeding were discussed.

MATERIAL AND METHODS

A mungbean cultivar, Chinese, was used as the experimental material. Two plants were grown in a 8-L pot filled with sandy soil in the experimental field of Kyushu University (33° 35' N, 130° 23' E). Seeds were sown on June 26 in 1997. A compound chemical fertilizer (N:P:K=16:16:16 in %) was applied 2 g/pot two weeks after germination of seeds. Full expanded young leaves excised from plants at the pre-flowering stage in the mid summer were used for measurements.

Measurement of CER

A leaf, the abaxial epidermis of which was peeled, is termed here PL, and an nonpeeled leaf is NL. CERs in PL and NL were measured at leaf temperatures of 15, 25 and 35°C in the air of different [CO₂] levels from 50 to 2,000 µmol mol⁻¹.

A closed system assimilation chamber (LD-2, Hansatech, UK) was partly remodeled and used here as an open system chamber to measure CER of a detached leaf (Fig. 1). The sampled air was monitored with an infrared CO₂ analyzer (LI-6262, LI-COR, USA). The air flow speed was 0.4 L min⁻¹ and the reading lag-time of the measurement system was 4 sec.

Water temperature in the water tanks equipped on the chamber (Fig. 1) was adjusted with a water temperature controller (CTER42WS, Komatsu-Yamato, Japan). A light of 2,000 µmol m⁻² s⁻¹ photosynthetically photon flux density (PPFD) was illuminated to a leaf disc (2.4 cm in diameter), and CER of the abaxial side in both PL and NL was measured. As a light source a slide projector (HILUX-HR, RIKAGAKU, Japan) with a 1 kw halogen lamp was used.

To prevent a leaf disk from drying, the air pumped into the chamber was moistened over 85% in relative humidity using a dew point controller (L-610, LI-COR, USA). The air sampled from the chamber was dehumidified by a dry agent (magnesium perchlorate) before it was sent into the CO₂ analyzer.

[CO₂] in the air was adjusted by mixing N₂, O₂ and CO₂ with a gas mixer (GM-3A, KOFLOC, Japan), and a CO₂ absorbent (soda lime) was used for the fine adjustment of [CO₂]. After the full openness of stomata was confirmed with NL, the CER was measured

in 50 to 2,000 $\mu\text{mol mol}^{-1}$ [CO₂]. The response of CER in PL was also measured at the same levels of [CO₂].

Simultaneous measurement of chlorophyll fluorescence and CER

The intensity of chlorophyll fluorescence at PSII was monitored with a portable fluorescence measurement system (MFMS-2T, Hansatech, UK). A leaf disc was fixed in the assimilation chamber (Fig. 1) and then placed in the dark for 30 min before fluorescence measurement in order to make the plastoquinone completely oxidized. After the dark treatment, an extremely weak measuring beam ($3.2 \mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD, 4.8 kHz) was applied to determine the initial fluorescence (F_0) of the leaf of unphotosynthetic situation. Then a 2 sec pulse illumination of $1,800 \mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD was given to read the maximum fluorescence (F_m).

After this, the leaf was redarkened for 2 min, and then a red actinic light ($300 \mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD) was continuously illuminated to let photosynthetic function start. Time courses of fluorescence (F_0) and CER were simultaneously monitored for 10 min, during which fluorescence peaks ($F_{x'}$) were measured by giving pulses of white saturation light at 20 sec intervals in the first 3 min and then at 30 sec intervals in the following 7 min. These were measured at 25 °C in the air of 2 and 20% [O₂]. The levels of [O₂] were adjusted with the gas mixer described above.

Based on measurements of fluorescence, non photochemical quenching (NPQ), quantum yield (Φ_e) at PSII and CO₂ quantum yield (Φ_{CO_2}) were calculated from the




Fig. 1. Diagram of the measurement system of CO₂ assimilation and chlorophyll fluorescence.

equations (1), (2) and (3), respectively.

$$\text{NPQ} = (F_m - F_m') / F_m' \quad \dots\dots\dots (1)$$

$$\Phi_e = (F_m' - F_o) / F_m' \quad \dots\dots\dots (2)$$

$$\Phi_{\text{CO}_2} = \text{CER} / (L \times i) \quad \dots\dots\dots (3)$$

where L is a light intensity supplied to a leaf disk, $L=300 \mu\text{molm}^{-2}\text{s}^{-1}$ PPFD is used here; i is the ratio of the photon absorbed by chlorophyll to the incident photon, $i=0.8$ is used.

The equation (4) represents the relationship between CER and Φ_e (Genty *et al.*).

$$\text{CER} = \Phi_e \times 0.5 \times L \times i / k \quad \dots\dots\dots (4)$$

Assuming the photon is evenly distributed to the two photosystems, 0.5 is used in the equation (4). The parameter k is the number of electron equivalents required to reduce 1mol CO_2 and calculated from the equation (5).

$$k = \Phi_e \times 0.5 \times L \times i / \text{CER} \quad \dots\dots\dots (5)$$

RESULTS AND DISCUSSION

$[\text{CO}_2]$ -CER response curves in PL and NL detached from plants grown in the mid summer were determined at 15, 25 and 35 °C, and shown in Fig. 2. CERs of both leaves at 35 °C linearly increased with $[\text{CO}_2]$ from 50 to 350 μmolmol^{-1} ; however, at a higher level of $[\text{CO}_2]$ than 350 μmolmol^{-1} CER of PL was higher than that of NL, and had a highest plateau of 26.0 to 26.8 $\mu\text{molm}^{-2}\text{s}^{-1}$ and 21.0 to 21.5 $\mu\text{molm}^{-2}\text{s}^{-1}$ in 500 to 1,000 μmolmol^{-1} $[\text{CO}_2]$ in PL and NL, respectively. Islam *et al.* (1994) suggested that maintaining stomatal aperture was the primary step toward the improvement of CER of mungbean cultivars. Also our results demonstrate that if the stomatal resistance is removed, the CER significantly increases with an increase in $[\text{CO}_2]$. But at a higher level above 1,000 μmolmol^{-1} $[\text{CO}_2]$, CER slightly decreased. As a reason for this, it is considered that the reproduction rate of substrate ribulose biphosphate (RuBP) becomes a limiting factor to photosynthesis, and also the excessive supply of substrate CO_2 is likely one of the inhibitors to the mesophyll activity.

CER decreased with decreasing leaf temperature from 35 to 15 °C (Fig. 2). The peaks of CER in NL appeared at a $[\text{CO}_2]$ of 500 μmolmol^{-1} , showing 18.2 and 13.4 $\mu\text{molm}^{-2}\text{s}^{-1}$ at 25 and 15 °C, respectively, and those of PL were 22.0 and 13.7 $\mu\text{molm}^{-2}\text{s}^{-1}$, respectively. The results in Fig. 2 indicate that the epidermis peeling was less effective on CER at 25 °C and not effective at 15 °C. At low temperatures the leaves had a peak of CER at a lower level of $[\text{CO}_2]$, the fact of which suggests that enzymes, such as Rubisco, related to CO_2 assimilation are inactivated at low temperatures so that the mesophyll photosynthetic activity may be saturated at a comparatively low $[\text{CO}_2]$ level.




Fig. 2. Responses of CER to $[\text{CO}_2]$ in PL (open) and NL (closed) measured at different leaf temperatures in a light intensity of $2,000 \mu\text{mol m}^{-2} \text{s}^{-1}$ PPFD.

CER in C_3 plants is obtained as a balanced value between CO_2 assimilation by RuBPCarboxylase (RuBPCase) and CO_2 release (photorespiration) by RuBPOxygenase (RuBPOase). The intercellular CO_2 concentration (C_i) of leaf adjusts the relative activity of carboxylase and oxygenase, by which CER and electron transport situation are changed. C_i varies with stomatal resistance.

In Table 1, CER, NPQ, Φ_e , Φ_{CO_2} and k at 35°C are compared between PL and NL. All these parameters were shown as averages during the photosynthetically steady state. CER of NL, $4.39 \mu\text{mol m}^{-2} \text{s}^{-1}$, was 45.5% of that ($9.65 \mu\text{mol m}^{-2} \text{s}^{-1}$) of PL, but Φ_e was 16.3% higher in NL (0.487) than in PL (0.419). The calculated value of $k=13.31$ in NL was about

Table 1. Effects of the leaf epidermis peeling on CER, NPQ, Φ_{CO_2} , Φ_e , and k. The measurement condition; $350 \mu\text{molmol}^{-1}$ $[\text{CO}_2]$, 20% $[\text{O}_2]$, $300 \mu\text{molmol}^{-2}\text{s}^{-1}$ PPFD and 35°C .

	CER ($\mu\text{molmol}^{-2}\text{s}^{-1}$)	NPQ	Φ_{CO_2}	Φ_e	k
NL	4.39 (45.5)	0.470 (136.2)	0.018 (45.0)	0.487 (116.3)	13.31 (255.5)
PL	9.65	0.345	0.040	0.419	5.21

PL, peeled leaf; NL, nonpeeled leaf; (), % of NL/PL

2.6-fold higher than that (5.21) of PL. This may be caused by the additional energy requirement by the increase in photorespiration rate in NL. NPQ (0.470) in NL was 36.2% higher than that of PL, the fact of which may suggest that a relatively large portion of the total energy was dispersed as heat or nonphotochemical quenching in NL because the CO_2 assimilation was restricted by stomatal resistance.

The value, $k=5.21$ shown in PL, is not so different from the theoretical minimum of $k=4$. This may suggest that the epidermis peeling is effective in promoting gas exchange and reducing photorespiration. If C_i comes close to the ambient $[\text{CO}_2]$ level of $350 \mu\text{molmol}^{-1}$ by removing the stomatal resistance, the CO_2 fixation is presumed to reach a nearly saturating level as predicted from Fig. 2.

Fig. 3 (A, B and C) represents the time courses of CER, NPQ and Φ_e measured with PL at 25°C in 20 and 2% $[\text{O}_2]$. CER quickly increased with time at the initial phase (Fig. 3A). Kobza and Edwards (1987) reported that RuBPCarboxylase–Oxygenase (Rubisco) in *Triticum aestivum* L. was light-activated within 1 min but the time required for reaching the steady state of photosynthesis was more than several minutes. It was also confirmed that the initial slope of CER in PL had a close relation with the light activation of Rubisco (Kubota et al., 1994). In Fig. 3A the initial slope in 2% $[\text{O}_2]$ indicates the light activation of RuBPCase, whereas that in 20% $[\text{O}_2]$ is shown as a joined value of RuBPCase and RuBPOase activations. The time required to reach the full activation was different in 2 and 20% $[\text{O}_2]$; it was less than 200 sec in the former and about 300 sec in the latter. Saturating values of CER in 2 and 20% $[\text{O}_2]$ were 12.50 and $7.88 \mu\text{molmol}^{-2}\text{s}^{-1}$, respectively (Fig. 3A), and both NPQ and Φ_e were a little higher in 20% $[\text{O}_2]$ (Fig. 3B, C).

CER, NPQ, Φ_e , Φ_{CO_2} and k in PL measured in 2 and 20% $[\text{O}_2]$ are shown in Table 2. CERs in 20 and 2% $[\text{O}_2]$ are termed CER_{20%} and CER_{2%} here, respectively. CER_{20%} of $7.88 \mu\text{molmol}^{-2}\text{s}^{-1}$ shown in Table 1 was 63.0% of $12.50 \mu\text{molmol}^{-2}\text{s}^{-1}$ of CER_{2%}. Contrary to this, the value of k, 7.75, in 20% $[\text{O}_2]$ was significantly higher than that (4.62) in 2% $[\text{O}_2]$. The photorespiration in 20% $[\text{O}_2]$ is considered as one of the causes of increasing the value of k, and if photorespiration is completely restricted, the theoretical minimum of k is 4. The value $k=4.62$ obtained here in 2% $[\text{O}_2]$ is 16% higher than the theoretical one.

Nine mol adenosine triphosphate (ATP) is consumed in the Calvin cycle in which 1 mol CO_2 is fixed by RuBPCase to produce 2 mol 3-phosphoglycerate (PGA) at the first step and 1 mol RuBP is reproduced at the final step. On the other hand, 1 mol CO_2 is




Fig. 3. Time courses of CER (A), NPQ (B) and Φ_e (C) in PL determined in the air of 2% [O₂] (open) and 20% [O₂] (closed). The other measurement conditions; 25 °C, 350 μmol mol⁻¹ [CO₂] and 300 μmol m⁻² s⁻¹ PPFD.

Table 2. CER, NPQ, Φ_{CO_2} , Φ_e and k of PL determined in 2% $[\text{O}_2]$ and 20% $[\text{O}_2]$. The measurement condition; $350 \mu\text{molmol}^{-1}$ $[\text{CO}_2]$, $300 \mu\text{molm}^{-2}\text{s}^{-1}$ PPFD and 25°C .

	CER ($\mu\text{molm}^{-2}\text{s}^{-1}$)	NPQ	Φ_{CO_2}	Φ_e	k
20% $[\text{O}_2]$	7.88 (63.0)	0.361 (103.4)	0.33 (63.5)	0.509 (105.9)	7.75 (167.8)
2% $[\text{O}_2]$	12.50	0.349	0.052	0.481	4.62
(); % of 20% $[\text{O}_2]$ / 2% $[\text{O}_2]$					

released by photorespiration after 3 mol PGA has been produced by RuBPOase, and reproduction of 1.5 mol RuBP from this 3 mol PGA requires 13.5 mol ATP in chloroplasts (Oliver, 1998). In the biochemical reaction of fixing 1 mol CO_2 , $k=4$ is required theoretically, and 9 mol ATP is consumed. On the other hand, the photorespiration (1 mol CO_2 release) consumes 13.5 mol ATP which is 1.5 times that required for 1 mol CO_2 fixation. If k has a parallel relation with the amount of ATP, $k=6$ ($=4 \times 1.5$) is given for photorespiration.

The photorespirationless CER determined in 2% $[\text{O}_2]$ is expressed as the equation (6).

$$\text{CER}_{2\%} = \Phi_{e2\%} \times L_0 / 4 \quad \dots\dots\dots (6)$$

where $\Phi_{e2\%}$ is a value of Φ_e determined in 2% $[\text{O}_2]$, and $L_0=0.5 \times L \times i$.

In the natural air condition where the photorespiration is performed, CER is expressed as the equation (7).

$$\text{CER} = \Phi_e \times L_1 / 4 - \Phi_e \times L_2 / 6 \quad \dots\dots\dots (7)$$

$$L_0 = L_1 + L_2 \quad \dots\dots\dots (8)$$

where L_0 is shared between L_1 and L_2 for CO_2 fixation and photorespiration, respectively.

Although the experimental value of $k=4.62$ (Table 2) is a little larger than the theoretical $k=4$, assuming here that $k=4.62$ is the fundamental value to fix 1 mol CO_2 , a value of k required for photorespiration is predicted as 6.93 (4.62×1.5) or 6.62 ($4 \times 1.5 + 0.62$). When $k=6.93$ is used, the equation (8) is rewritten as the equation (9).

$$\text{CER} = \Phi_e \times L_1 / 4.62 - \Phi_e \times L_2 / 6.93 \quad \dots\dots\dots (9)$$

By substituting CER and Φ_e shown in Table 1 and 2 into the equation (9), and $L_0=120$ ($=300 \times 0.5 \times 0.8$) $\mu\text{molm}^{-2}\text{s}^{-1}$ into the equation (8), the total CO_2 fixation rate ($\Phi_e \times L_1 / 4.62$), photorespiration rate ($\Phi_e \times L_2 / 6.93$) and the photorespiration ratio (Photorespiration rate/Total CO_2 fixation rate) are calculated and listed up in Table 3. The highest photorespiration rate and ratio, $3.30 \mu\text{molm}^{-2}\text{s}^{-1}$ and 42.9%, are estimated in NL in 20%

Table 3. Estimated rates of photorespiration and total fixed CO₂ of PL based on the data listed in Table 1 and 2.

	[CO ₂] ($\mu\text{mol mol}^{-1}$)	[O ₂] (%)	°C	Φ_e	k	CER ($\mu\text{mol m}^{-2}\text{s}^{-1}$)	TC ($\mu\text{mol m}^{-2}\text{s}^{-1}$)	PR ($\mu\text{mol m}^{-2}\text{s}^{-1}$)	PR/TC (%)
NL	350	20	35	0.487	13.31	4.39	7.70	3.30	42.9
PL	350	20	35	0.419	5.21	9.65	10.14	0.49	4.9
PL	350	20	25	0.509	7.75	7.88	10.02	2.14	21.3
PL	350	2	25	0.481	4.62	12.50	12.50	0	0

TC, Total CO₂ fixation rate; PR, Photorespiration rate, PR/TC; Photorespiration ratio.

[O₂]. Reduction in these values of PL in 20% [O₂] is considered to depend on the increase in Ci by epidermis peeling. The photorespiration ratio in PL measured at 35 °C in 20% [O₂] is 4.9%, which is a considerably low compared to that (21.3%) in PL at 25 °C. The epidermis peeling promotes gas exchange and hence the photorespiration is restricted as mentioned above. However, as shown in table 3, the strength of the restriction is considerably different among the individual measurements of PL. As a reason for this it may be considered that the relative humidity in the assimilation chamber, though being kept over 85% here, was not controlled so rigorously, so that change in air moisture could affect the evapotranspiration and Ci.

The photorespiration in leaf is estimated from the electron transport according to the method described above, but a further examination is necessary to determine more adequate values of k and also to make more rigorous the environmental control of the measurement system. The cyclic electron transport from PSI and Mehler reaction are not taken into account here for the sake of simplicity. Also because it may be considered that these functions are less operative in our experimental condition where the light intensity used is not so high (300 $\mu\text{mol m}^{-2}\text{s}^{-1}$ PPFD).

In general, photorespiration rate is estimated by subtracting CER_{20%} from CER_{2%}. However the calculated in this manner is not thought to represent the real rate of photorespiration in 20% [O₂]. The calculation, as tried here, based on electron transport may be considered to have a better consistency to the real photorespiration rate than that obtained by the subtraction (CER_{2%} - CER_{20%}). The obtained values are expected to be a useful indicator for evaluating the photorespiratory traits among genotypes.

To maintain stomatal aperture and raise mesophyll activity are essential for improving photosynthesis of leaves grown in adequate growth environments. But such a C₃ plant as mungbean grown in the tropics frequently closes stomata by being subjected to water deficits in strong solar radiation and high temperature. If CO₂ assimilation is restricted in high light intensities, the chemical energy produced by photosystem is accumulated in chloroplasts, which becomes a cause of damaging the photosynthetic apparatus. Under the condition like this, dispersing the accumulated energy is an indispensable role of photorespiration.

Cultivars having a high photorespiratory ability is considered to disperse effectively excessive energy in leaf, the function of which allows leaves to retard senescence and maintain a photosynthetic activity for a longer time. However, having a high photores-

piration rate concurrently means a large production loss by CO₂ release. In order to utilize the genetic variation in photorespiratory characteristics as an effective selection criterion in breeding, it is necessary to deepen the study on the relative strength of its positive and negative effects on mungbean production in relation to the growth conditions.

REFERENCES

- Genty, B., J.-M. Briantais and N. Baker 1989 The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence. *Biochim. Biophys. Acta.*, **990**: 87–92
- Harbinson, J., B. Genty and N. R. Baker 1990 The relationship between CO₂ assimilation and electron transport in leaves. *Photosynth. Res.*, **25**: 213–224
- Islam, M. T., W. Agata and F. Kubota 1994 Evaluation of characteristics related to photosynthetic rate and leaf area production in mungbean genotypes. *Jpn. J. Crop Sci.*, **63**: 502–509
- Kobza, J. and G. E. Edwards 1987 The photosynthetic induction response in wheat leaves: net CO₂ uptake, enzyme activation, and leaf metabolites. *Planta*, **171**: 549–559
- Krall, J. P. and G. E. Edwards 1990 Quantum yields of of photosystem II electron transport and carbon dioxide fixation in C₃ plants. *Aust. J. Plant Physiol.*, **17**: 579–588
- Krall, J. P. and G. E. Edwards 1991a Environmental effects on the relationship between the quantum yields of carbon assimilation and *in vivo* PSII electron transport in maize. *Aust. J. Plant Physiol.*, **18**: 267–278
- Krall, J. P., G. E. Edwards and M. S. B. Ku 1991b Quantum yield of photosystem II and efficiency of CO₂ fixation in *Flaveria* (Asteraceae) species under varying light and CO₂. *Aust. J. Plant Physiol.*, **18**: 369–383
- Kubota, F., W. Agata and M. Morokuma 1991 Photosynthetic control factors in a single leaf of sweet potato, *Ipomoea batatas* L. 1. Leaf epidermis peeling method and photosynthetic rate of peeled leaf. *Jpn. J. Crop Sci.*, **60**: 510–514
- Kubota, F., M. Yatomi and W. Agata 1992 CO₂ exchange rate in sweet potato (*Ipomoea batatas* Lam.) leaves with without lower epidermis. *Photosynthetica*, **26**: 257–260
- Kubota, F., N. Nada and W. Agata 1994 Photosynthetic control factors in a single leaf of sweet potato, *Ipomoea batatas* Lam. 3. Estimation of *in vivo* rubisco activity from the CO₂ exchange rate of a peeled leaf. *Jpn. J. Crop Sci.*, **63**: 89–95
- Marco G. Di, F. Manes, D. Tricoli and E. Vitale 1990 Fluorescence parameters measured concurrently with net photosynthesis to investigate chloroplastic CO₂ concentration in leaves of *Quercus ilex* L. *J. Plant Physiol.*, **136**: 538–543
- Oliver D. J. 1998 Photorespiration and the C₂ cycle. In "Photosynthesis: A comprehensive treatise" ed. by A. S. Raghavendra, Cambridge Univ. Press, Cambridge, pp. 173–182
- Shreiber, U., U. Schliwa and W. Bilger 1986 Continuous recording of photochemical and non-photochemical chlorophyll fluorescence quenching with a new type of modulation fluorometer. *Photosynth. Res.*, **10**: 51–62
- Shreiber, U., W. Bilger, H. Hormann and C. Neubauer 1998 Chlorophyll fluorescence as a diagnostic tool: basics and some aspects of practical relevance. In "Photosynthesis: A comprehensive treatise" ed. by A. S. Raghavendra, Cambridge Univ. Press, Cambridge, pp. 320–336